

Rahil Malik MD, Fellow of American College of Obstetrics and Gynecology

1200 N University Drive, Plantation, Florida, 33322 Tel: 954-791-3090 Fax: 954-791-3166 www.DavieObGyn.com

IUD PLACEMENT CONSENT FORM FOR

I hereby request and authorize Dr Malik to perform the procedure: IUD placement

Description: The IUD is one of the most effective birth control methods available. Fewer than 1/100 people per year using the IUD will get pregnant. Possible **Advantages** of using the IUD include: Highly effective and long lasting contraception, Low risk of side effects, Reversible (immediate return to fertility after removal), Decreased menstrual blood loss (Hormonal IUDs), Good safety record, High satisfaction rates

Potential side effects/risks of IUD use: Mild to moderate pain when the IUD is inserted, Cramping or backache for a few days after insertion, 2-10% of IUD users spontaneously expel their IUD within the first year of use, Irregular or absent menses (Mirena, Kyleena and Skyla), Heavier, longer and/or crampier menstrual periods (Paragard IUD) which may improve with time. Spotting between menses which usually, improves or resolves within 3 months (both Hormonal IUDs and Paragard IUD), Anemia (only if heavy menses occurs, Serious but very infrequent risks of IUD use: Infection of the uterus or tubes {pelvic inflammatory disease (PID)} within the first 3 weeks after the IUD is inserted. If PID is not treated, infertility could result, While pregnancy with an IUD in place is rare, if you are pregnant you are at increased risk of an ectopic pregnancy (pregnancy outside of the uterus) which can be life threatening if not treated, Perforation of the uterine wall during IUD placement or embedment in the uterine wall (rare, surgery may be required), At the time of insertion, there is a possibility that the cervix will be tightly closed and not allow for insertion of the IUD into the uterus. Another visit may need to be scheduled. You may be asked to return while on your period or after taking a medication to soften your cervix to improve success of insertion.

The IUD does not protect against HIV (the virus that causes AIDS) or other sexually transmitted infection (STIs). Using a condom correctly and consistently helps prevent STIs. Patient may elect to have the IUD removed at any time. A visit with a health care provider is needed to have the IUD removed. An alternative method of contraception should be used if the IUD strings cannot be located. Emergency care is always available if you should need it.

I have reviewed the Paragard, Mirena, Kyleena and Skyla Patient Information handouts. I have been informed of the IUD insertion and removal procedure and what to expect when the IUD is inserted and know I should check the strings regularly. I have reviewed and understand all of the above information. I have been given the opportunity to ask questions and have had them answered to my satisfaction. After reviewing the above information, I hereby authorize and direct my doctor to insert the IUD.

Patient Signature	Date
	 Date
Dr Malik Signature	Date

